

## **General Disclaimer**

### **One or more of the Following Statements may affect this Document**

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

"Made available under NASA sponsorship  
in the interest of early and wide dis-  
semination of Earth Resources Survey  
Program information and without charge  
for any use made thereon."

RECEIVED  
NASA STI FACILITY  
ACQ. BR.

F MAY 28 1976 Exch

DCAF# 1019452  
1 2 3 4 5

**Title of Investigation:** 28990 Investigation of Environmental  
Change Pattern in JAPAN

E7.6-10360  
CR-14795-4

**Principal Investigator:** Takakazu MARUYASU  
Science University of Tokyo  
Noda City, Chiba ken , 278, JAPAN

**Quarterly Progress Report for Period**

January 1976-March 1976

Original photography may be purchased from:  
EROS Data Center  
10th and Dakota Avenue  
Sioux Falls, SD 57198

**Name and address of**

**National Sponcering**

**Agency**

Science and Technology Agency  
Kasumigaseki 2-2-1 Chiyoda ku,  
Tokyo, 100, JAPAN

**Subject**

1. Investigation of Soil Erosion in Hokkaido which is caused by  
Thawing of Soil Water in Late Spring

**Co-Investigator:** Shigechika HAYASHI

Upland Farming Division,  
National Agricultural Experiment  
Station,  
Memuro, Hokkaido, 082, JAPAN

(E76-10360) INVESTIGATION OF ENVIRONMENTAL  
CHANGE PATTERN IN JAPAN. INVESTIGATION OF  
SOIL EROSION IN HOKKAIDO WHICH IS CAUSED BY  
THAWING OF SOIL WATER IN LATE SPRING  
QUARTERLY PROGRESS REPORT, (SCIENCE UNIV.

N76-25607  
HC \$3.50

UNCLAS  
00360


G3/43

Distribution of density response was determined about two scenes, the one obtained on 11 June 1975, the other, 24 May 1973, in the same test area (figure-1). Results are showed in figure-2.

The reflectance of 6, 7-bands is larger in the imagery of June than that of May. This fact is attributed to the increased crop coverage due to the seasonal difference between May and June, because in this district in Hokkaido, only wheat and grass show considerable coverage in May, and wheat grows not so large in this season yet, on the other hand, in June, wheat grows to cover nearly 100% of the soil surface and sugar-beet also grows to show about 40% coverage.

Each of three soil types (Alluvial soil, Andosol-Brown and Andosol-Black) has its own characteristic reflectance response among four bands in the imagery of May. This relationship is also recognized in the imagery of June, when the soils are already covered by crops to some extent.

The rectangular shapes of individual fields where wheat or grass were planted are distinguished clearly by looking over these enlarged imageries of June. Therefore, it may be possible to observe the status of crops more in detail by enlarging these imageries to a still larger scale.



(a) Band 5 image



(b) Band 7 image

Fig.1 Selected test site (A: Alluvial soil, B: Andosol-Brown, C: Andosol-Black)





Fig.2 Distribution of density response for three different soil type field within four bands of B & W bulk positives. (24 May 1973, 11 June 1975)